

The South Lancashire Flora
(Vascular Plants Family 045 Vitaceae)


Vitales

VITACEAE

Grape-vine family

Vitis vinifera L.

Grape-vine



Status & distribution: Archaeophyte-cultivated; naturalised near habitation.


Earliest record traced: Climbing over bushes by a road at Formby (SD20Y, Liverpool Botanical Society, 1942).

Habitat: Hedgerows, roadsides, on the site of old sewage farms, on waste ground and tips.

Comments: Native to mainland Europe.

Parthenocissus quinquefolia (L.) Planchon

Virginia-creeper



Status & distribution: Neophyte-naturalised; locally naturalised usually in or nearby to built up areas.


Earliest record traced: Haresfinch Bank (SJ5296, Anon@Biobank, 1983).

Habitat: Waste ground, old walls, hedges, roadsides and scrub near houses.

Comments: Native to North America.

Parthenocissus inserta (A. Kernr) Fritsch

False Virginia-creeper



Formby 2009

Status & distribution: Neophyte-naturalised; widely semi-established usually in or nearby to built up areas.


Earliest record traced: Manchester, Bolton & Bury Canal, Salford (SJ8399, DP Earl, 1992).

Habitat: Waste ground, hedgerows, old walls, former mills, tips, and in parks.

Comments: Native to North America.

Parthenocissus tricuspidata (Seibold & Zucc.) Planchon

Boston Ivy





Formby Powerhouse DP Earl 2009

Status & distribution: Neophyte-surviving; spreading to the Leeds & Liverpool Canal towpath at Bootle (SJ3491, J Clitherow & H Ash, 1983) and (SJ3394 & SJ3493 DP & J Earl); Formby Powerhouse (SD20X, PH Smith & MP Wilcox, 2004).

Habitat: Derelict grounds, walls and planted areas.

Comments: Native to East Asia.